

HOW TO USE

DO NOT apply oil sprays within 10 days of applying chlorothalonil.

PLANTS	DISEASE	RATE	HOW TO APPLY
Tomatoes, Capsicums, Cucumbers, Strawberries, Zucchini	Powdery Mildew	30-40 g per 10 L of water and add 20 mL emulsified vegetable oil	Begin application at first sign of disease. Repeat application at 7-10 day intervals. Use the higher rate and shorter repeat interval under conditions of rapid plant growth or severe disease pressure. Apply to ensure complete and thorough coverage of all plant surfaces. Pay particular attention to the underside of leaves.
Roses	Powdery Mildew Black Spot		
Geraniums (pelargoniums)	Rust (<i>Puccinia pelargonii-zonalis</i>)	40 g per 10 L of water and add 30 mL emulsified vegetable oil	Begin application at first sign of disease. Repeat application at 7 day intervals. Apply to ensure complete and thorough coverage of all plant surfaces. Pay particular attention to the underside of leaves.
Grapevines	Powdery Mildew (<i>Oidium</i> spp.)	Mix 40 g per 10 L of water and add 20 mL emulsified vegetable oil	Begin application at first sign of disease. Repeat applications at intervals of 10 to 14 days as new growth or infection occurs, or if conditions favour disease. Apply to ensure complete and thorough coverage of foliage.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED.
WITHOLDING PERIOD (WHP): NOT REQUIRED WHEN USED AS DIRECTED.



Organic Crop Protectants

READ SAFETY DIRECTIONS



eco-fungicide®

HOME GARDEN FUNGICIDE

ACTIVE CONSTITUENT: 940 g/kg POTASSIUM BICARBONATE

FOR THE MANAGEMENT OF FUNGUS DISEASES IN ROSES, GRAPEVINES, GERANIUMS, VEGETABLES & STRAWBERRIES IN THE HOME GARDEN AS PER DIRECTIONS FOR USE.

NET CONTENTS
500 g



MAKES
UP TO 160 L
OF SPRAY

HOW TO PREPARE

eco-fungicide is a contact fungicide containing activated potassium bicarbonate that requires direct contact with the fungus organism for control. **eco-fungicide** disrupts the fungus cell, causing the cell walls to collapse. However, under conditions highly favourable to the disease, control may not be achieved and suppression only can be expected.

eco-fungicide rapidly kills both powdery mildew spores and hyphae.

MIXING: Half fill spray tank. While stirring, slowly add required amount of **eco-fungicide** and then the required amount of emulsified vegetable oil e.g. **eco-oil**. Top up with required quantity of water. Mix well before commencing spray procedure. Do not allow mixtures to stand. Use within 24 hours of mixing.

APPLICATION: Contact with the disease organism is essential for control. Apply in sufficient water to ensure complete and thorough coverage of all plant surfaces including the undersides of leaves. Ensure plants are not suffering from any environmental stress such as moisture stress or heat stress when applying **eco-fungicide** to control high levels of powdery mildew or black spot infection as water loss from the infection site can cause death of plant tissue.

CAUTION: DO NOT allow chemical containers or spray to get into drains, sewers, streams or ponds.

STORAGE AND DISPOSAL: Store in the closed, original container in a cool, dry place out of the reach of children. Do not store in direct sunlight. Dispose of empty container by wrapping in paper, placing in plastic bag and putting in garbage.

SAFETY DIRECTIONS: May irritate the eyes. Avoid contact with eyes. Wash hands after use.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126.

FOR EMERGENCIES ONLY CALL: 1800 033 111 (24 HOURS).

eco-fungicide and **eco-oil** are registered trade marks and the OCP drop device is a trade mark of Organic Crop Protectants.

ORGANIC CROP PROTECTANTS

1956 Dandenong Rd, Clayton VIC 3168
Telephone: 1800 634 204

For further information visit
www.ecoorganicgarden.com.au

APVMA Approval No. 64510/57313

EAN BARCODE
9336099000326